



המחלקה
לפיזיותרפיה

אוניברסיטת
אריאל
בשומרון

נספח לקווים מנחים לקווים מנחים להורדת סיכון להתפתחות תופעות לוואי ברביע עליון, אחרי ניתוחים וטיפולים בסרטן שד (2020)

פרופ' תמר יעקב, PT, MPT, PhD, ראש התוכנית לתואר שני בפיזיותרפיה, אוני' אריאל בשומרון.

גב' גיליאן ברכה, MScPT, מדריכה מוסמכת לפי שיטת Casley Smith, פיזיותרפיסטית לימפטית, קליניקה פרטית, קיבוץ מעגן מיכאל.

גב' רות פלג, MScPT, מדריכה מוסמכת לפי שיטת Casley Smith, פיזיותרפיסטית אחראית במערך אונקולוגיה-פלסטיקה, מרכז שרות לטיפול בלימפדמה, מרכז רפואי ת"א ע"ש סוראסקי.

גב' עמית אפשטיין, BPT, Graduate Comprehensive Oncology Rehabilitation (GCOR), פיזיותרפיסטית לימפטית, מובילה ארצית לתחום האונקולוגיה בפיזיותרפיה במכבי שירותי בריאות.

גב' אילנה יושע, פיזיותרפיסטית לימפטית, עמותת לימפדמה בישראל.

מקורות

1. Leclerc AF, Jerusalem G, Devos M, Crielaard JM, Maquet D. Multidisciplinary management of breast cancer. Arch Public Health. 2016;74:50.
2. Hayes SC, Johansson K, Stout NL, et al. Upper-body morbidity after breast cancer: incidence and evidence for evaluation, prevention, and management within a prospective surveillance model of care. Cancer. 2012;15:118(8 Suppl):2237-2249.
3. Ellegaard MB, Grau C, Zachariae R, Jensen AB. Women with breast cancer report substantially more disease- and treatment-related side or late effects than registered by clinical oncologists: a cross-sectional study of a standard follow-up program in an oncological department. Breast Cancer Res Treat. 2017;164 (3):727-736.
4. Mehnert A, Koch U. Psychological comorbidity and health-related quality of life and its association with awareness, utilization, and need for psychosocial support in a cancer register-based sample of long-term breast cancer survivors. J Psychosom Res. 2008;64 (4):383-391.
5. Armer JM, Hulett JM, Bermas M, Ostby P, Stewart BR, Cormier JN. Best practice guidelines in assessment, risk reduction, management, and surveillance for post-breast cancer lymphedema. Curr breast Cancer Rep. 2013;5(2):134-144.
6. Moffatt C, Doherty D, Morgan P. Best Practice for the Management of Lymphedema- International consensus. 2006. <https://www.lympho.org/portfolio/best-practice-for-the-management-of-lymphoedema/>. Accessed: 13.8.19.
7. Gho SA, Steele JR, Jones SC, Munro BJ. Self-reported side effects of breast cancer treatment: a cross-sectional study of incidence, associations, and the influence of exercise. Cancer Causes Control. 2013;24(3):517-528.
8. Torre Lacombe M, del Moral OM, Coperias Zazo JL, Yuste Sánchez MJ, Ferrandez JC, Zapico Goñi A. Axillary Web Syndrome after axillary dissection in breast cancer: a prospective study. Breast Cancer Res Treat. 2009;117(3):625-630.
9. Jacob T, Bracha J. Identification of signs and symptoms of Axillary Web Syndrome and breast seroma during a course of physical therapy, seven months post lumpectomy: A case report. Phys Ther. 2019;99(2):229-239.
10. Tuschy B, Berlitz S, Romero S, et al. Clinical aspects of intraoperative radiotherapy in early breast cancer: short-term complications after IORT in women treated with low energy x-rays. Radiat Oncol. 2013;8:95. doi: 10.1186/1748-717X-8-95.
11. Falco M, Masojc B, Rollaa M, et al. Risk factors for seroma evacuation in breast cancer patients treated with intraoperative radiotherapy. Rep Pract Oncol Radiother. 2016;21:225-231.
12. Ribeiro Pereira ACP, Koifman RJ, Bergmann A. Incidence and risk factors of lymphedema after breast cancer treatment: 10 years of follow-up. Breast. 2017 Oct 6; 36:67-73.

13. Palesh O, Scheiber C, Kesler S, Mustian K, Koopman C, Schapira L. Management of side effects during and post-treatment in breast cancer survivors. *Breast J*. 2018;24:167-175.
14. Nielsen I, Gordon S, Selby A. Breast cancer-related lymphedema risk reduction advice: a challenge for health professionals. *Cancer Treat Rev*. 2008; 34(7):621-628.
15. Rogan S, Taeymans J, Luginbuehl H, Aebi M, Mahnig S, Gebruers N. Therapy modalities to reduce lymphedema in female breast cancer patients: a systematic review and meta-analysis. *Breast Cancer Res Treat*. 2016;159:1-14.
16. Torre Lacomba M, Mariña Jose M, Sañchez Y, et al. Effectiveness of early physiotherapy to prevent lymphedema after surgery for breast cancer: randomized, single blinded, clinical trial. *BMJ*. 2010; 340:b5396. doi: 10.1136/bmj.b5396.
17. Box RC, Reul-Hirche HM, Bullock-Saxton JE, Furnival CM. Shoulder movement after breast cancer surgery: results of a randomized controlled study of post-operative physiotherapy. *Breast Cancer Res Treat*. 2002;50:35-50.
18. Stout Gergich NL, Pfalzer LA, McGarvey C, Springer B, Gerber LH, Soballe P. Preoperative assessment enables the early diagnosis and successful treatment of lymphedema. *Cancer*. 2008;112(12): 2809-2819.
19. Schmitz KH, Troxel AB, Andrea Cheville A, et al. Physical Activity and Lymphedema (The PAL Trial): Assessing the safety of progressive strength training in breast cancer survivors. *Contemp Clin Trials*. 2009; 30(3):233-245.
20. Friedenreich CM, Neilson HK, Megan S, Courneya KS. Physical Activity and Cancer Outcomes: A Precision Medicine Approach. *Clin Cancer Res*. 2016;22(19):4766-4775.
21. Ahn S, Port ER. Lymphedema Precautions: Time to Abandon Old Practices? *J Clin Oncol*. 2016;34(7):655-658.
22. Ferguson CM, Swaroop MN, Horick N, et al. Impact of ipsilateral blood draws, injections, blood pressure measurements, and at travel on the risk of lymphedema for patients treated for breast cancer. *J Clin Oncol*. 2016;34:691-698.
23. Ochalek K, Gradalski T, Partsch H. Preventing early postoperative arm swelling and lymphedema manifestation by compression sleeves after axillary lymph node interventions in breast cancer patients: A randomized controlled trial. *J Pain Symptom Manage*. 2017;54(3):346-354.
24. Shiang-Ru Lu, Rong-Bin Hong, Willy Chou, and Pei-Chi Hsiao. Role of physiotherapy and patient instruction in lymphedema control following breast cancer surgery. *Ther Clin Risk Manag*. 2015;11:319-327.
25. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *CA Cancer J Clin*. 2016;66(1):43-73.
26. Box RC, Reul-Hirche HM, Bullock-Saxton JE, Furnival CM. Breast Physiotherapy after breast cancer surgery: a RCT to minimize lymphedema. *Breast Cancer Res Treat*. 2002;64:75:51-64.
27. Hanna L, Grothaus K, Peterson ML, Bertram S. Breast cancer survivors' perceptions of the delivery of lymphedema education in the United States. *J Womens Health Phys Therap*. 2017;41(3):137-144.
28. Meade E, McIlpatrick S, Groarke AM, Butler E, Dowling M. Survivorship care for postmenopausal breast cancer patients in Ireland: What do women want? *Eur J Oncol Nurs*. 2017;28:69-76.
29. Keesing S, Rosenwax L, McNamara B. A call to action: The need for improved service coordination during early survivorship for women with breast cancer and partners. *Women & Health*. 2019;59(4): 406-419.
30. Stout NL, Binkley JM, Schmitz KH, et al. A prospective surveillance model rehabilitation for women with breast cancer. *Cancer* 2012; 118 (8 suppl) 2191-200.
31. Lymphedema. Clalit HMO.
https://www.clalit.co.il/he/medical/medical_diagnosis/Pages/lymphedema.aspx Accessed June 26 2019. [Hebrew].
32. Maccabi Physical Therapy Department. Maccabi HMO.:
<https://www.maccabi4u.co.il/15079-he/Maccabi.aspx?TabId=24285>. Accessed June 26 2019 [Hebrew].
- Meuhedet HMO. Patients' rights. <https://www.meuhedet.co.il/> . Accessed June 26 2019 [Hebrew].
33. Thangaratinam S, Redman C. The Delphi technique. *The Obstetrician & Gynecologist* 2005;7:120-125.
34. Decision-making for guideline development at WHO, chapter 16, p; 206-7. In: WHO handbook for guideline development- 2nd ed, 2014. http://www.who.int/publications/guidelines/Chp16_May2016.pdf. Accessed June 26 2019.

35. de Villiers MR, de Villiers PTJ, & Kent AP. The Delphi technique in health sciences instruction research. *Med Teach*. 2005;27(7); 639-643.
36. Burns PB, Rohrich RJ, Chung KC. The Levels of Evidence and their role in Evidence-Based Medicine. *Plast Reconstr Surg*. 2011;128(1):305-310.
37. Szajewska H. Evidence-Based Medicine and Clinical Research: Both Are Needed, Neither Is Perfect. *Ann Nutr Metab*. 2018; (suppl 3):13-23.
38. Nutly DD. The adequacy of response rates to online and paper surveys: what can be done? *Assess Eval High Educ*. 2008;33(3):301-314.
39. Jacob T, Bracha J. Lymphedema treatment certified physical therapists' awareness of and practice patterns for side effect risk reduction after breast cancer treatments. *Int J Physiother*. 2019;6(5):224-230.
40. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008. 336(7650):924-926.
41. Guyatt GH, Oxman AD, Kunz R, et al. Going from evidence to recommendation. *BMJ*. 2008 336(7650):1049-1051.
43. Cemal Y, Pusic A, and Mehrara BJ. Preventative measures for lymphedema: Separating fact from fiction. *J Am Coll Surg*. 2011;213(4):543-551.
44. Nudelman J. Debunking Lymphedema Risk-Reduction Behaviors: Risky Conclusions. *Lymphat Res Biol*. 2016;14(3):124-126.
45. Nudelman J. Do No Harm: Lymphedema risk reduction behaviors. *J Clin Oncol*. 2016; 34(25):3109-3110.
46. Devoogdt N, Geraerts I, Van Kampen M, et al. Manual lymph drainage may not have a preventive effect on the development of breast cancer-related lymphoedema in the long term: a randomised trial. *J Physiother*. 2018;64(4):245-254.
47. Johansson K, Tribe K, Weibull A, Newton RC. Low intensity resistance exercise for breast cancer patients with arm lymphedema with or without compression sleeve. *Lymphology* 2005;38(4):167-180.
48. Partsch H, Stout N, Forner-Cordero I, et al. Clinical trials needed to evaluate compression therapy in breast cancer related lymphedema (BCRL). Proposals from an expert group. *Int Angiol*. 2010;29(5):442-253.
49. Todd J, Scally A, Dodwell D, Horgan K, Topping A. A randomised controlled trial of two programs of shoulder exercise following axillary node dissection for invasive breast cancer. *Physiotherapy*. 2008;94(4):265-273.
50. Singh BRU, Newton P, Cormie DA, et al. Effects of compression on lymphedema during resistance exercises in women with breast cancer-related lymphedema: A randomized, cross-over trial. *Lymphology*. 2015;48:80-92.
51. Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine. American College of Sports Medicine Roundtable on Exercise Guidelines for Cancer. *Med Sci Sports Exerc*. 2010;42(7):1409-1426.
52. Rafn BS, Hung S, Hoens AM, et al. Prospective surveillance and targeted physiotherapy for arm morbidity after breast cancer surgery: a pilot randomized controlled trial. *Clin Rehabil*. 2018;32(6):811-826.
53. Kilbreath SL, Refshauge KM, Beith JM, et al. Upper limb progressive resistance training and stretching exercises following surgery for early breast cancer: a randomized controlled trial. *Breast Cancer Res Treat*. 2012;133(2):667-676.
54. Nelson NL. Breast cancer related lymphedema and resistance exercises: A systematic review. *Strength Con Res*. 2016; 30(9):2656-2665.
55. Furmaniak AC, Menig M, Markes MH. Exercise for women receiving adjuvant therapy for breast cancer (Review). *Cochrane Database of Systematic Reviews* 2016;9. Art. No.: CD005001.
56. van der Leeden M, Huijsmans RJ, Geleijn E, et al. Tailoring exercise interventions to comorbidities and treatment-induced adverse effects in patients with early stage breast cancer undergoing chemotherapy: a framework to support clinical decisions. *Disabil and Rehab*. 2008; 40 (4): 486-496.
57. Lynch BM, Neilson HK, Friedenreich CM. Physical activity and breast cancer prevention .Recent Results. *Cancer Res*. 2011;186:13-42.
58. Khan F, Amatya B, Pallant JF, Rajapaksa I, Brand C. Multidisciplinary rehabilitation in women following breast cancer treatment: a randomized controlled trial. *J Rehabil Med*. 2012;44(9):788-794.

59. Kim J, Choi WJ, Jeong SH. The effects of physical activity on breast cancer survivors after diagnosis. *J Cancer Prev.* 2013;18(3):193-200.
60. Benjaminsen Borch K, Braaten T, Lund E, Weiderpass E. Physical activity before and after breast cancer diagnosis and survival - the Norwegian women and cancer cohort study. *BMC Cancer.* 2015;15:967. doi: 10.1186/s12885-015-1971-9.
61. Hilfiker R, Meichtery A, Eicher M, et al. Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: a systematic review incorporating an indirect-comparisons meta-analysis. *Br J Sports Med.* 2018;52(10):651-658.
62. Espíndula RC, Nadas GB, Rosa MID, de Araujo FC, Grande AJ. Pilates for breast cancer: A systematic review and meta-analysis. *Rev Assoc Med Bras.* 2017; 63 (11):1006-1012.
63. Pan Y, Yang K, Wang Y, Zhang L, Liang H.. Could yoga practice improve treatment-related side effects and quality of life for women with breast cancer? A systematic review and meta-analysis. *Asia Pac J Clin Oncol.* 2017;13 (2):e79-e95.
64. Sprod LK, Hsieh CC, Hayward R, and Schneider CM. Three versus six months of exercise training in breast cancer survivors. *Breast Cancer Res Treat.* 2010;121(2):413-419.
65. Jönsson C, Johansson K. Pole walking for patients with breast cancer-related arm lymphedema. *Physiotherapy Theory Pract.* 2009;25(3):165-173.
66. Di Blasio A, Morano T, Napolitano G, et al. Nordic walking and the Isa method for breast cancer survivors: Effects on upper limb circumferences and total body extracellular water - a Pilot Study. *Breast Care (Basel).* 2016;11(6):428-431.
67. Shaw C, Mortimer P, Judd PA. A randomized controlled trial of weight reduction as a treatment for breast cancer-related lymphedema. *Cancer.* 2007;110(8):1868-1874.
68. Pekmezi DW, Demark-Wahnefried W. Updated evidence in support of diet and exercise interventions in cancer survivors. *Acta Oncol.* 2011;50(2):167-178.
69. Dieli-Conwright CM, Lee K, Kiwata JL. Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. *Curr Breast Cancer Rep.* 2016;8(3):139-150.
70. Kootstra JJ, Dijkstra PU, Rietman JS, Hoekstra-Weebers J. A longitudinal study of shoulder and arm morbidity in breast cancer survivors 7 years after sentinel lymph node biopsy or axillary lymph node dissection. *Breast Cancer Res Treat.* 2013;139:125-134.
71. Tsuchiya M, Masujima M, Kato T, et al. Knowledge, fatigue, and cognitive factors as predictors of lymphedema risk-reduction behaviors in women with cancer. *Support Care Cancer.* 2019;27(2):547-555.
72. Boccardo FM1, Ansaldo F, Bellini C et al. Prospective evaluation of a prevention protocol for lymphedema following surgery for breast cancer. *Lymphology.* 2009;42(1):1-9.
73. McNeely ML, Campbell K, Ospina M, et al. Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database Syst Rev.* 2010;16;(6):CD005211. doi: 10.1002/14651858.CD005211.pub2.
74. Springer BA, Levy E, McGarvey C, et al. Pre-operative assessment enables early diagnosis and recovery of shoulder function in patients with breast cancer. *Breast Cancer Res Treat.* 2010;120(1):135-147.
75. Stuiver MM, ten Tusscher MR, Agasi-Idenburg CS, Lucas C, Aaronson NK, Bossuyt PM.. Conservative interventions for preventing clinically detectable upper-limb lymphoedema in patients who are at risk of developing lymphoedema after breast cancer therapy. *Cochrane Database Syst Rev.* 2015;13;(2):CD009765. doi: 10.1002/14651858.CD009765.pub2. 2015(2).
76. De Groef A, Van Kampen M, Dieltjens E, et al. Effectiveness of postoperative physical therapy for upper-limb impairments after breast cancer treatment: a systematic review. *Arch Phys Med Rehabil.* 2015;96(6):1140-1153.
77. Binkley JM, Harris SR, Levangie PK, et al. Patient perspectives on breast cancer treatment side effects and the prospective surveillance model for physical rehabilitation for women with breast cancer. *Cancer.* 2012;15;118(8 Suppl):2207-2216.
78. Taylor C, Shewbridge A, Harris J, Green JS. Benefits of multidisciplinary teamwork in the management of breast cancer. *Breast Cancer.* 2013;5:79-85.
79. Casley-Smith. *Modern Treatment for Lymphoedema.* 5th ed. The Lymphoedema Association of Australia, Inc. 1997;126-127.

80. Zuther JE, Norton S. Lymphedema Management. The Comprehensive Guide for Practitioners 4th ed. Thieme Medical Publishers, Inc. 2018; 559.
81. Foldi M, Foldi E, Kubic S (eds). Textbook of Lymphology for Physicians and Lymphedema Therapists. 1st ed. Urban and Fischer Verlag 2003.
82. NLN Position Papers. <https://lymphnet.org/position-papers>. Accessed 1.7.2019.
83. Epstein RM and Street RL. The values and value of patient-centered care. *Ann Fam Med*. 2011;9(2) (Editorial).
84. Rinder SH. Pretreatment lymphedema education and identified educational resources in breast cancer patients. *Patient Educ Couns*. 2006; 61(1):72-79.
85. Ostby PL, Armer, JM, Smith K, and Stewart BR. Patient Perceptions of Barriers to Self-Management of Breast Cancer-Related Lymphedema. *West J Nurs Res*. 2018;40(12):1800–1817.
86. Playdon M, Ferrucci LM, McCorkle R, et al. Health information needs and preferences in relation to survivorship care plans of long-term cancer survivors in the American Cancer Society's Study of Cancer Survivors-I. *J Cancer Surviv*. 2016;10(4):674–685.